

AD-A034 659

EDGEWOOD ARSENAL ABERDEEN PROVING GROUND MD  
THE EFFECT OF INTRAVENOUS THIAMINE HYDROCHLORIDE ON PRALIDOXIME--ETC(U)  
DEC 76 J JOSSELSO, F R SIDELL  
EB-TR-76115

F/G 15/2

UNCLASSIFIED

NL


| OF |

AD  
A034659



END

DATE  
FILMED  
2-77



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ADA034659

12  
b.s.

AD

EDGEWOOD ARSENAL TECHNICAL REPORT

EB-TR-76115

THE EFFECT OF INTRAVENOUS THIAMINE HYDROCHLORIDE  
ON PRALIDOXIME PHARMACOKINETICS IN MAN

by

John Josselson, M.D., MAJ, MC  
Frederick R. Sidell, M.D.

Biomedical Laboratory

December 1976

D D C  
RECEIVED  
JAN 21 1977  
REGISTERED  
C



DEPARTMENT OF THE ARMY  
Headquarters, Edgewood Arsenal  
Aberdeen Proving Ground, Maryland 21010



Approved for public release; distribution unlimited.

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER EB-TR-76115	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE EFFECT OF INTRAVENOUS THIAMINE HYDROCHLORIDE ON PRALIDOXIME PHARMACOKINETICS IN MAN.		5. TYPE OF REPORT & PERIOD COVERED Technical Report. Aug 74 - Sep 74
6. AUTHOR(s) John Josselson M.D., MAJ, MC Frederick R. Sidell M.D.		7. PERFORMING ORG. REPORT NUMBER
8. PERFORMING ORGANIZATION NAME AND ADDRESS Commander, Edgewood Arsenal Attn: SAREA-BL-O Aberdeen Proving Ground, MD 21010		9. CONTRACT OR GRANT NUMBER(s)
10. CONTROLLING OFFICE NAME AND ADDRESS Commander, Edgewood Arsenal Attn: SAREA-TS-R Aberdeen Proving Ground, MD 21010		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1W7627AD2502
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. REPORT DATE December 1976
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15. NUMBER OF PAGES 17
15. SECURITY CLASS. (of this report) UNCLASSIFIED		16. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Medical Defense Against Chemical Agents, Prophylaxis and Therapy for Lethal Agents		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Pralidoxime chloride      Renal excretion Thiamine hydrochloride      Plasma concentration Human subjects		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Six subjects were given pralidoxime chloride (2-PAMCl) 5 mg/kg, iv, by itself and during a 2.5-hour intravenous infusion of thiamine hydrochloride (100 mg/hr); pralidoxime was given one hour after the infusion was started and the infusion ran 1.5 hours after pralidoxime was given. In the subjects receiving thiamine the plasma concentration of oxime was greater throughout the test period, the renal excretions and clearance of oxime were less during the first 1.5 hours, and the half-life of oxime was prolonged. These data suggest that thiamine inhibits the excretion of oxime, thus prolonging its effects. Whether this will be of value in the therapy of anticholinesterase compound intoxication is unknown.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED 401007  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) LB

## PREFACE

The work described in this report was authorized under Project/Task No. 1W762710AD25-02, Medical Defense Against Chemical Agents, Prophylaxis and Therapy for Lethal Agents. This work was started in August 1974 and completed in September 1974.

The volunteers in these tests are enlisted US Army personnel. These tests were governed by the principles, policies, and rules for medical volunteers as established in AR 70-25 and the Declaration of Helsinki.

The use of trade names in this report does not constitute an official endorsement or approval of the use of such commercial hardware or software. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, Edgewood Arsenal, Attn: SAREA-TS-R, Aberdeen Proving Ground, Maryland 21010; however Defense Documentation Center and the National Technical Information Service are authorized to reproduce the document for US Government purposes.

The stamp is tilted and contains the following text and markings:

- At the top right, there is a checkmark.
- Below the checkmark, the text "RECEIVED" is visible.
- Below "RECEIVED", the text "DISTRIBUTION" is visible.
- Below "DISTRIBUTION", the text "BY" is visible.
- Below "BY", the text "DISTRIBUTION AVAILABILITY CODES" is visible.
- Below "DISTRIBUTION AVAILABILITY CODES", the text "BY" is visible.
- Below "BY", the text "AVAIL. and/or SPECIAL" is visible.
- At the bottom left, there is a signature that appears to be "A".

CONTENTS

	<u>Page</u>
I. INTRODUCTION . . . . .	5
II. EXPERIMENTAL . . . . .	5
A. Subjects . . . . .	5
B. Design . . . . .	5
C. Controls . . . . .	6
D. Thiamine Treatment . . . . .	6
E. Calculations . . . . .	6
F. Complications . . . . .	7
III. RESULTS . . . . .	7
A. Clinical Findings . . . . .	7
B. Plasma Concentration . . . . .	7
C. Urinary Excretion of Pralidoxime . . . . .	7
D. Renal and Plasma Clearance . . . . .	8
IV. KINETIC DATA . . . . .	8
A. Volumes of Distribution . . . . .	8
B. Plasma Half-Life . . . . .	12
C. Equilibrium Constants . . . . .	12
V. DISCUSSION . . . . .	12
LITERATURE CITED . . . . .	14
DISTRIBUTION LIST . . . . .	15



## THE EFFECT OF INTRAVENOUS THIAMINE HYDROCHLORIDE ON PRALIDOXIME PHARMACOKINETICS IN MAN

### I. INTRODUCTION

Pralidoxime chloride (pyridine-2-aldoxime methochloride)\*, an adjunct to atropine in the treatment of anticholinesterase intoxication, is rapidly and almost completely (80-90%) excreted in the urine<sup>1-6</sup>. This necessitates frequent administration in those circumstances in which a prolonged plasma concentration is desirable.

Swartz *et al.*,<sup>5</sup> recently reported that concurrent intramuscular administration of thiamine hydrochloride with intravenous pralidoxime prolongs plasma half-life and delays the renal elimination of oxime. This suggests that pralidoxime might be handled by the kidney as a strong organic base and might be secreted by the same mechanism by which other strong bases (e.g., thiamine, choline, guanidine, tetraethylammonium) are secreted. Thus, concurrent administration of two bases competing for the same mechanism might lead to delayed secretion of one or both<sup>7</sup>.

Those results led us to explore further the clinical use of thiamine hydrochloride to potentiate plasma oxime levels. This report describes the pharmacokinetics of pralidoxime when administered one hour after the start of a continuous infusion of thiamine hydrochloride.

### II. EXPERIMENTAL

#### A. Subjects.

The subjects were US Army enlisted men who agreed to the protocol after thorough explanation and discussion. Screening included complete physical and laboratory examinations\*\*. A detailed history of allergy was obtained and intradermal scratch testing with thiamine was performed to exclude potentially hypersensitive subjects. All scratch tests were negative; two potential subjects were excluded from the study on the basis of a strongly atopic history.

#### B. Design.

Six subjects participated and received pralidoxime with and without thiamine, each serving as his own control. Subjects entered the ward on the evening before the study, ate a light breakfast about one hour before the study began, and drank about 1 to 2 liters of fluid in the two hours prior to testing. Each subject was tested at one-week intervals in a randomized crossover design plan.

---

\*Protopam<sup>R</sup>, Ayerst Laboratories, New York, NY 10017.

\*\*Chest X-ray, ECG, complete blood count, routine urinalysis, blood urea nitrogen, serum creatinine, liver function tests (SGOT, alkaline phosphatase, serum bilirubin).



### C. Controls.

At the start of testing, a continuous infusion of normal saline into the antecubital vein was begun and continued for 2.5 hours. One hour after the infusion was started, each subject received an intravenous injection of 5 mg/kg of pralidoxime chloride over two minutes through the scalp needle of the infusion line.

### D. Thiamine Treatment.

At the start of testing, a continuous infusion containing normal saline and thiamine hydrochloride\* was begun; thiamine was delivered at a rate of 100 mg/hr over 2.5 hours (2.0 mg thiamine/ml normal saline, 50 ml/hr). One hour after the infusion was begun, each subject received an iv injection of pralidoxime, 5 mg/kg, as in the control trial.

Urine from each subject was pooled for the following periods: 0-1.5, 1.5-3, 3-6, and 6-24 hours. Urine was collected for an additional 12 hours from four thiamine-treated subjects. Plasma specimens were obtained through an indwelling scalp needle in a vein in the opposite arm, kept open with heparin, at 0.05, 0.10, 0.15, 0.25, 0.50, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, and 4.0 hours after the injection of oxime. Plasma and urine were analyzed by the method of Groff *et al.*<sup>8</sup> for oxime. Pulse and blood pressure, and subjective responses to both thiamine and pralidoxime, were closely monitored. Subjects were kept in bed during the 2.5-hour infusion; then they were allowed to walk around.

### E. Calculations.

The serial plasma oxime concentrations plotted against time fit the curve described by the bi-exponential equation<sup>1</sup>

$$C = Ae^{-\alpha t} + Be^{-\beta t} \quad (1)$$

which describes the disposition of drug in a two-compartment model. In this equation, C is the plasma concentration, t is the time, A and B represent the zero-time intercepts of the linear components of the rapid and slow portions of the plasma decay curve, and  $\alpha$  and  $\beta$  represent the slopes of these two lines divided by -2.303. A more detailed description of the derivation of this equation is provided by Wagner<sup>9</sup> and discussed with respect to pralidoxime in studies by Swartz *et al.*<sup>4,5</sup> and Sidell<sup>1</sup>. The computer program NONLIN<sup>10</sup> was used to estimate the parameters of this equation. From them the following were calculated:

1. The half-life of the rapid equilibrium phase ( $t_{1/2\gamma}$ ) and the slower elimination phase ( $t_{1/2\beta}$ ).
2. The volumes of distribution of the central compartment ( $V_1$ ), the peripheral compartment ( $V_2$ ), and the total volume ( $V_d$ ).

\*Thiamine Hydrochloride Injection U.S.P. 10 mg/cc, Natcon Chemical Co., Inc., Plainview, NY.

3. The rate constants for drug movement from central to peripheral compartment ( $K_{12}$ ), from peripheral to central compartment ( $K_{21}$ ), and for elimination from the central compartment ( $K_{13}$ ).

4. The plasma and renal clearances of pralidoxime.

The plasma clearance was calculated by dividing the dose of pralidoxime by the area under the plasma concentration versus time curve (which was obtained from the integral of equation 1); the renal clearance was obtained by dividing the amount of pralidoxime excreted into the urine during a given time period by the area under the curve of that time interval.

#### F. Complications.

Because of difficulties encountered during administration of oxime to two thiamine-treated subjects, rapid intravenous infusion was not achieved; hence, plasma data on only four thiamine-treated subjects were used for calculations of the parameters listed above compared to control data from six subjects.

Likewise, loss of one urine specimen (1.5-3 hours) from one control subject prevented calculating each parameter.

### III. RESULTS

#### A. Clinical Findings.

Both thiamine and pralidoxime were well tolerated by all subjects. No objective or subjective changes were noted after initiation of thiamine infusion. Following pralidoxime, all subjects complained of mild-to-moderate visual symptoms characterized as "blurring" or visual heaviness, lasting five to 10 minutes. Tests of extraocular muscle movement, accommodation, and light reflex were normal in all men. No changes in blood pressure or pulse rate were noted following oxime injection. Intravenous injection of thiamine did not potentiate any of the subjective experiences and did not alter pulse rate and blood pressure after oxime was given.

#### B. Plasma Concentration (Six Control Subjects; Four Thiamine-Treated Subjects).

The mean plasma concentrations of pralidoxime are plotted in the following figure. After six minutes, the plasma concentrations of the thiamine-treated group were significantly higher than those of the control group ( $p < 0.05$ ; paired t test) at each time interval during the remaining hours of blood sampling.

#### C. Urinary Excretion of Pralidoxime.

Over the 24-hour period of urine collection, there was no significant difference in total oxime excretion between the control group and thiamine-pretreated groups. The former excreted 85.7% ( $\pm 6.2$  S.D.,  $n=5$ ) and the latter 87.2% ( $\pm 5.2$ ,  $n=6$ ) of the injected pralidoxime. However, in the first 1.5 hours the thiamine-treated men eliminated significantly less oxime

(38.2% vs 72.4% of the total 24-hour urinary oxime excretion,  $p < 0.001$ , paired t test,  $n=5$ ) than the subjects in the control group and, in the remaining 22.5 hours, significantly more (46.3% vs 12.8%,  $p < 0.01$ , paired t test,  $n=5$ ) (table 1).




Figure. Mean Plasma Concentrations of Pralidoxime

#### D. Renal and Plasma Clearance.

Mean renal and plasma clearance data are presented in table 2. The renal clearance of oxime at 1.5 hours was significantly reduced by the simultaneous administration of thiamine ( $p < 0.01$ , paired t test,  $n=4$ ), and this maintained over 24 hours. Similarly, the plasma clearance of pralidoxime was greatly reduced by the administration of thiamine ( $p < 0.01$ , paired t test,  $n=4$ ).

### IV. KINETIC DATA

The kinetic data are shown in table 3.

#### A. Volumes of Distribution.

Following thiamine administration, there was a definite decrease in the total volume of distribution ( $p < 0.05$ , paired t test,  $n=4$ ), characterized by a large decrease in the peripheral (extravascular) volume,  $V_2$  ( $p < 0.05$ ), and a moderate increase in the central compartment ( $p < 0.05$ ).



Table 1. Urinary Recovery of Pralidoxime

(a) Mean Cumulative Excretion of Pralidoxime Over 24 Hours\*

	Control	$\bar{P}$ Thiamine	P
0-1.5 hr	72.4 $\pm$ 3.3	38.2 $\pm$ 9.0	<.001
0-3 hr	87.2 $\pm$ 2.4	53.7 $\pm$ 4.2	<.01
0-6 hr	97.0 $\pm$ 1.0	75.9 $\pm$ 8.6	<.01
0-24 hr	100	100	—

\* Expressed as percent of the total amount of pralidoxime excreted in the urine  $\pm$ S.D.

(b) Mean Urinary Excretion of Pralidoxime for Each Collection Period

	Control	$\bar{P}$ Thiamine	P
0-1.5 hr	72.4 $\pm$ 3.3	38.2 $\pm$ 9.0	<.001
1.5-3 hr	14.8 $\pm$ 2.4	15.5 $\pm$ 2.1	>.70
3-6 hr	9.8 $\pm$ 2.1	22.2 $\pm$ 2.2	<.01
6-24 hr	3.0 $\pm$ 1.0	24.1 $\pm$ 8.6	<.01
Total	100	100	—



Table 2. Renal and Plasma Clearances (ml/min)

Case number	Renal clearance at 1.5 hours		Renal clearance at 24 hours		Plasma clearance	
	c*	t**	c*	t**	c*	t**
6731	477	-	437	-	548	-
6748	502	-	527	-	631	-
6736	546	109	535	253	565	293
6717	515	132	496	216	536	286
6718	574	245	-	441	726	489
6730	717	283	848	350	741	385
Mean	555	205	529	315	625	363
S.D. $\pm$	86.23	97.86	77.06	101.2	90.71	95.20
p <sup>†</sup>	<0.01		<0.001		<0.01	
n	4		3		4	

\* Oxime alone

\*\* Oxime and Thiamine

† Paired t test

Table 3. Pharmacokinetic Constants

Case	A ( $\mu\text{g/ml}$ )	$\alpha$ ( $\text{hr}^{-1}$ )	B ( $\mu\text{g/ml}$ )	e ( $\text{hr}^{-1}$ )	$t_{1/2\alpha}$ (hr)	$t_{1/2\beta}$ (hr)	$K_{21}$ ( $\text{hr}^{-1}$ )	$K_{13}$ ( $\text{hr}^{-1}$ )	$K_{12}$ ( $\text{hr}^{-1}$ )	$V_1$ (ml/kg)	$V_2$ (ml/kg)	$V_D$ (ml/kg)
<u>CONTROL</u>												
6731	29.0	9.8	5.4	0.53	0.07	1.30	1.99	2.61	5.69	145	415	560
6748	21.2	11.9	5.8	0.79	0.059	0.88	3.15	2.96	6.53	185	385	570
6736	23.3	8.3	4.6	0.51	0.083	1.25	1.83	2.52	4.52	179	443	622
6717	26.8	10.1	8.1	0.66	0.069	1.05	2.83	2.34	5.56	143	281	425
6718	19.3	9.6	4.7	0.61	0.072	1.14	2.38	2.47	5.38	208	470	678
6730	17.5	10.1	4.4	0.66	0.069	1.05	2.56	2.59	5.57	228	496	
Mean	22.9	10.0	5.5	0.63	0.070	1.11	2.46	2.56	5.54	182	415	597
S.D. $\pm$	4.4	1.1	1.4	0.09	0.01	0.15	0.50	0.21	0.64	33.7	76.5	105
<u>THIAMINE-TREATED</u>												
6731												
6748												
6736	9.8	3.8	8.7	0.47	0.18	1.46	2.08	0.86	1.36	268	175	443
6717	14.0	6.1	10.8	0.42	0.11	1.64	2.87	0.89	2.71	202	191	393
6718	12.4	4.1	5.5	0.47	0.17	1.46	1.59	1.23	1.79	280	319	594
6730	15.7	4.8	5.4	0.42	0.14	1.63	1.53	1.32	2.35	238	364	601
Mean	13.0	4.7	7.6	0.45	0.15	1.55	2.02	1.08	2.05	247	261	508
S.D. $\pm$	2.5	1.0	2.7	0.03	0.03	0.10	0.62	0.23	0.60	34.7	0.924	105.6
p*	0.05	0.001	N3	0.05	0.01	0.02	N3	0.001	0.001	0.05	0.05	0.05

\* By paired t test N = 4

## B. Plasma Half-Life.

Following thiamine administration, significant increases occurred in both the short half-time,  $t_{1/2\gamma}$  (0.07 hr vs 0.15 hr,  $p < 0.01$ ,  $n=4$ ) and, more importantly, the long half-life of elimination,  $t_{1/2\beta}$  (1.11 hr vs 1.55 hr,  $p < 0.02$ ,  $n=4$ ).

## C. Equilibrium Constants.

Likewise, following thiamine, highly significant decreases occurred in the rate constants  $K_{12}$  ( $p < 0.001$ ,  $n=4$ ) and  $K_{13}$  ( $p < 0.001$ ,  $n=4$ ). The change in  $K_{12}$  closely parallels the decrease in the peripheral compartment noted above.

## V. DISCUSSION

These data indicate that concurrent intravenous administration of thiamine hydrochloride significantly decreases both the renal and plasma clearance and delays the renal elimination of pralidoxime chloride apparently without altering the mode of excretion. Such treatment was associated with substantially higher plasma concentrations of oxime than in the controls throughout the 4-hour period of blood sampling. This study extends the investigation of Swartz *et al.*<sup>5</sup> (who administered thiamine intramuscularly with the oxime) by intravenous pretreatment with the vitamin to achieve and maintain maximal plasma levels at the time of oxime administration.

A dose of 100 mg of thiamine per hour was empirically selected for the pretreatment dose. Neither pharmacokinetic data nor renal tubular secretory transport maxima are available for thiamine, and it is possible that still larger doses of the vitamin would further retard the renal elimination of oxime.

During the 1.5 hours of thiamine administration after injection of pralidoxime, the amount of oxime excreted in the urine was halved. This suggests that prolonging the thiamine infusion at the same rate might have maintained this difference for a longer period of time.

The fractional recovery of oxime in the urine, the plasma half-life for elimination, and the renal and plasma clearances of the drug noted are comparable to those noted previously<sup>1,5</sup>. Swartz<sup>5</sup>, however, reported no significant changes in volumes of distribution after intramuscular thiamine administration, though a tendency did appear toward a reduction in the peripheral compartment ( $V_2$ )\*. In contrast, our data show significant changes in all those volumes (table 3). A definite explanation for this phenomenon is not possible, but the alternatives include: (1) increased protein binding affinity for oxime following thiamine administration (but since pralidoxime is negligibly bound, this seems unlikely); (2) increased sequestration of pralidoxime by other organs (e.g., CNS, liver); and (3) alteration of erythrocytic and/or vascular membrane permeabilities by as yet unknown mechanisms.

\*Personal communication.



While this study does not offer proof, these data suggest that pralidoxime may be excreted, at least in part, by a renal secretory mechanism shared by several other strong organic bases<sup>7</sup> although this does not exclude the possibility that several different mechanisms may be involved in the renal elimination of oxime. Ultimate identification of the responsible mechanism will depend upon further studies with specific blockers of secretion and reabsorption as well as the use of micropuncture techniques.

The effect of thiamine in prolonging the plasma half-life of pralidoxime has been demonstrated, although the clinical value of this regimen in the therapy of anticholinesterase poisoning has not been investigated. Thiamine appears well tolerated in pharmacologic doses and has an extremely low incidence of undesirable side effects; however, a few case reports indicate that a rare but fatal allergic reaction to parenteral (particularly intravenous) administration of thiamine may occur and is presumably anaphylactic<sup>11-13</sup>.

These results suggest that larger doses of thiamine might produce even larger increases in the half-life of pralidoxime, and it is possible that other strong organic bases with quaternary  $N^+$  sites – such as guanidine, choline, and tetraethylammonium – might be equally effective. Additional studies are currently under way.



## LITERATURE CITED

1. Sidell, F. R., Groff, W. A., and Kaminskis, A. Toxogonin and Pralidoxime: Kinetic Comparison After Intravenous Administration to Man. *J. Pharm. Sci.*, **61**, 1765-9 (November 1972).
2. Kondritzer, A. A., Zvirblis, P., Goodman, A., and Paplanus, S. H. Blood Plasma Levels and Elimination of Salts of 2-PAM in Man After Oral Administration. *J. Pharm. Sci.*, **57**, 1142-6, (July 1968).
3. Loomis, T. A. Distribution and Excretion of Pyridine-2-aldoxime Methiodide (PAM); Atropine and PAM in Sarin Poisoning. *Toxicol. and Appl Pharmacol.*, **5**, 489-99 (1963).
4. Swartz, R. D., and Sidell, F. R. Effects of Heat and Exercise on the Elimination of Pralidoxime in Man. *Clin. Pharm. Ther.*, **14**, 83-89 (1973).
5. Swartz, R. D., and Sidell, F. R. Renal Tubular Secretion of Pralidoxime in Man. Edgewood Arsenal Technical Report EB-TR-73017. January 1974. UNCLASSIFIED report.
6. Berglund, F., Elwin, D. E., and Sundwall, A. Studies on the Renal Elimination of N-methyl-pyridinium-2-aldoxime. *Biochem. Pharmacol.*, **2**, 383-388 (1962).
7. Peters, L. Renal Tubular Excretion of Organic Bases. *Pharmacol. Rev.*, **12**, 1-35 (1960).
8. Groff, W. A., and Ellin, R. J. A New and Rapid Determination of Pyridinium Aldoximes in Blood and Urine. *Clin. Chem.*, **15**, 72-83 (1962).
9. Wagner, J. C. Pharmacokinetics. J. M. Richard Laboratories, Grosse Pointe, Michigan. 1969.
10. Metzler, C. M. A User's Manual for NONLIN. The Upjohn Company. November 25, 1969.
11. Smith, J. A., Fo, P. P., Weinstein, H. R., Ludwig, A. S., and Weitherion, J. M. Some Aspects of Thiamine Toxicity. *J. Pharm. Exp. Ther.*, **93**, 294-304 (July 1948).
12. Haley, T. J., and Flesher, A. M. A Toxicity Study of Thiamine Hydrochloride. *Science*, **104**, 567-8 (December 1946).
13. Reingold, J. M., and Webb, F. R. Sudden Death Following Intravenous Injection of Thiamine Hydrochloride. *J.A.M.A.*, **130**, 491-2 (1946).

# DISTRIBUTION LIST 4

Names	Copies	Names	Copies
EDGEWOOD ARSENAL		Director	
FOREIGN INTELLIGENCE OFFICER	1	Defense Intelligence Agency	
CHIEF, LEGAL OFFICE	1	Attn: DB-4G1	1
CHIEF, SAFETY OFFICE	1	Washington, DC 20301	
CHIEF, PLANS OFFICE		DEPARTMENT OF THE ARMY	
Attn: SAREA-PL	1		
CDR, US ARMY TECHNICAL ESCORT CENTER	1	HQDA (DAMO-SSC)	1
PUBLIC HEALTH SERVICE LO	1	WASH DC 20310	
AUTHOR'S COPY, Biomedical Laboratory	2		
BIOMEDICAL LABORATORY		Director	
Attn: SAREA-BL-M	7	Defense Civil Preparedness Agency	
Attn: SAREA-BL-O	1	Attn: RE	1
Attn: SAREA-BL-Q	1	Attn: PO(DC)	1
Attn: SAREA-BL-R	1	Washington, DC 20301	
Attn: SAREA-BL-RC	1		
Attn: SAREA-BL-RE	1	CINCUSAREUR	
Attn: SAREA-BL-RI	1	Attn: AEAGC-DS	1
Attn: SAREA-BL-V	1	APO New York 09403	
CHEMICAL LABORATORY			
Attn: SAREA-CL	1	Chief, Office of Research,	
Attn: SAREA-CL-B	1	Development & Acquisition	
Attn: SAREA-CL-BS	1	Attn: DAMA-CSM-CM	1
Attn: SAREA-CL-C	1	Attn: DAMA-ARZ-D	1
Attn: SAREA-CL-D	1	Washington, DC 20310	
Attn: SAREA-CL-P	1	Headquarters, Sixth US Army	
Attn: SAREA-CL-TE	1	NBC Office	
		Attn: AFKC-OP-PN	1
DEVELOPMENT AND ENGINEERING DIRECTORATE		Presidio of San Francisco, CA 94129	
Attn: SAREA-DE-D	1		
Attn: SAREA-DE-EP	1	HQDA (DAMI-FIT)	1
Attn: SAREA-DE-ET	1	WASH, DC 20310	
Attn: SAREA-DE-M	1		
Attn: SAREA-DE-N	1	CINCUSAREUR	
		Attn: AEAMD-PM	1
MANUFACTURING TECHNOLOGY DIRECTORATE		APO New York 09403	
Attn: SAREA-MT-D	1		
Attn: SAREA-MT-E	1	US ARMY HEALTH SERVICE COMMAND	
Attn: SAREA-MT-P	1		
DIRECTOR OF PRODUCT ASSURANCE		Commander	
Attn: SAREA-PA-A	1	US Army Environmental Hygiene Agency	
Attn: SAREA-PA-P	1	Attn: HSE-AA-L, Librarian, Bldg 2100	1
Attn: SAREA-PA-Q	1	APG-Edgewood Area	
DIRECTOR OF TECHNICAL SUPPORT			
Attn: SAREA-TS-R	2	Superintendent	
Attn: SAREA-TS-L	3	Academy of Health Sciences	
Attn: SAREA-TS-E	1	US Army	
		Attn: HSA-CDC	1
		Attn: HSA-IHE	1
		Fort Sam Houston, TX 78234	
DEPARTMENT OF DEFENSE			
Administrator		OFFICE OF THE SURGEON GENERAL	
Defense Documentation Center			
Attn: Accessions Division	12	HQDA (SGRD-HR)	1
Cameron Station		Forrestal Building	
Alexandria, VA 22314		WASH DC 20314	

# DISTRIBUTION LIST 4 (Contd)

Names	Copies	Names	Copies
US ARMY MATERIEL DEVELOPMENT AND READINESS COMMAND		Commander Pine Bluff Arsenal Attn: SARP-B-ETA Pine Bluff, AR 71611	1
Commander US Army Materiel Development and Readiness Command Attn: DRCDE-DM Attn: DRCLDC Attn: DRCMT Attn: DRCSF-S 5001 Eisenhower Ave Alexandria, VA 22333	1 1 1 1	US ARMY TRAINING & DOCTRINE COMMAND	
Commander US Army Foreign Science & Technology Center Attn: DRXST-ISI 220 Seventh St., NE Charlottesville, VA 22901	2	Commandant US Army Infantry School Combat Support & Maintenance Dept. Attn: NBC Division Fort Benning, GA 31905	1
Project Manager for Chemical Demilitarization and Installation Restoration Attn: DRCPM-DR, Bldg E4585 APG-Edgewood Area	1	Commandant US Army Missile & Munitions Center & School Attn: ATSK-CD-MD Attn: ATSK-DT-MU-EOD Redstone Arsenal, AL 35809	1 1
Redstone Scientific Information Center Attn: Chief, Documents US Army Missile Command Redstone Arsenal, AL 35809	1	Commandant US Army Military Police School Attn: ATZN-CD Fort McClellan, AL 36205	1
Director DARCOM Field Safety Activity Attn: DRXOS-C Charlestown, IN 47111	1	Chief, Training Literature Division Attn: ATZN-TDP Fort McClellan, AL 36205	1
US ARMY ARMAMENT COMMAND		Headquarters US Army Military Police School/Training Center Attn: ATSJ-TD-CR Fort McClellan, AL 36205	1
Commander US Army Armament Command Attn: DRSAR-ASN Attn: DRSAR-IMB-C Attn: DRSAR-RDM Attn: DRSAR-RDT Rock Island, IL 61201	1 1 1 1	Commander US Army Infantry Center Attn: ATZB-CD-MS-C Fort Benning, GA 31905	1
Commander US Army Dugway Proving Ground Attn: Technical Library, Docu Sect Dugway, UT 84022	1	Commandant US Army Ordnance Center & School Attn: ATSL-CD-MS Attn: ATSL-DTG APG-Aberdeen Area	1 1
Commander Rocky Mountain Arsenal Attn: SARRM-EA Attn: SARRM-MD Denver, CO 80240	1 1	US ARMY TEST & EVALUATION COMMAND	
		Record Copy CDR, APG Attn: STEAP-AD-R/RHA APG-Edgewood Area, Bldg E5179	1



## DISTRIBUTION LIST 4 (Contd)

Names	Copies	Names	Copies
CDR, APG Attn: STEAP-TL APG-Aberdeen Area	1	Chief, Bureau of Medicine & Surgery Department of the Navy Attn: CODE 5 Washington, DC 20372	1
Commander US Army Test & Evaluation Command Attn: DRSTE-FA APG-Aberdeen Area	1	Commander Naval Weapons Center Attn: A. B. Galloway/Code 3171 Attn: Technical Library/Code 233 China Lake, CA 93555	1 1
Commander US Army Cold Regions Test Center Attn: STECR-TD APO Seattle 98733	1	DEPARTMENT OF THE AIR FORCE  Air University Library Attn: AUL/LSE-8879 Maxwell AFB, AL 36112	1
Commander US Army Tropic Test Center Attn: STETC-MO-A (Tech Library) APO New York 09827	1	HQ Foreign Technology Division (AFSC) Attn: PDRR Wright-Patterson AFB, OH 45433	1
Commander Dugway Proving Ground Attn: STEDP-PO Dugway, UT 84022	1	Commander Aeronautical Systems Division Attn: ASD/AELD Wright-Patterson AFB, OH 45433	1
DEPARTMENT OF THE NAVY  Chief of Naval Research Attn: Code 443 800 N. Quincy Street Arlington, VA 22217	1	HQ, USAF/SGPR Forrestal Bldg WASH DC 20314	1
Commander Naval Facilities Engineering Command Attn: Code 03 200 Stovall Street Alexandria, VA 22332	1	HQ, USAF/RDPN WASH DC 20330	1
Commander Naval Explosive Ordnance Disposal Facility Attn: Army Chemical Office Indian Head, MD 20640	1	HQ, Ogden ALC/MMWRA Hill AFB, UT 84406	1
Commander Naval Surface Weapons Center Attn: Tech Lib & Info Svcs Br White Oak Laboratory Silver Spring, MD 20910	1	Director Air Force Inspection and Safety Center Attn: IGD(AFISC/SEV) Norton AFB, CA 92409	1
Commander Naval Surface Weapons Center Dahlgren Laboratory Attn: DX-21 Dahlgren, VA 22448	1	Commander Armament Development & Test Center Attn: DLOSL (Technical Library) Eglin AFB, FL 32542	1
Commander Nuclear Weapons Training Group, Atlantic Naval Air Station Attn: Code 21 Norfolk, VA 23511	1	NORAD Combat Operations Center/DBN Cheyenne Mtn Complex, CO 80914	1
		OUTSIDE AGENCIES  Battelle, Columbus Laboratories Attn: TACTEC 505 King Avenue Columbus, OH 43201	1
		Director of Toxicology National Research Council 2101 Constitution Ave, NW Washington, DC 20418	1